

## Supplementary Material

**Supplementary Table 1:** Association of CFT parameters with coronary tortuosity

	<b>Tortuosity <math>\geq</math> moderate (n=34; 38%), n (%)</b>	<b>Tortuosity &lt;moderate (n=55; 62%), n (%)</b>	<b>p-value</b>
<b>ACh testing</b>			
Epicardial spasm	5 (15%)	14 (25%)	0.29
Microvascular spasm	26 (76%)	28 (51%)	<b>0.02</b>
No spasm	3 (9%)	13 (24%)	0.09
<b>Microvascular function testing</b>			
CFR, median (Q1–Q3)	2.53 (2.05–3.05)	2.60 (2.19–3.06)	0.72
CFR $\geq$ 2.5	18 (53%)	33 (60%)	0.65
CFR <2.5	16 (47%)	22 (40%)	0.65
HMR, median (Q1–Q3)	1.75 (1.47–2.19)	1.63 (1.41–1.98)	0.18
HMR $\leq$ 2.5	30 (89%)	53 (96%)	0.19
HMR >2.5	4 (12%)	2 (4%)	0.19

**Supplementary Figure 1:** Diagnostic Yield of Invasive Coronary Functional Testing and Distribution of Functional Coronary Disease Endotypes in Patients with ANOCA; CMD = Coronary Microvascular Dysfunction

