

# Supplemental Material

Supplemental Table 1. Sensitivity Analyses using the win ratio method .....2

### Supplemental Table 1. Sensitivity Analyses using the win ratio method

<b>Sensitivity analysis strategy</b>	<b>Win ratio</b>	<b>95% CI</b>	<b>P-value</b>
Replacing cardiovascular death with all-cause mortality	1.18	1.03 — 1.36	0.0209
Excluding time to first hospitalization for arrhythmias from the hierarchy;	1.22	1.05 — 1.42	0.0106
Analyzing the change in KCCQ-OSS at 12 instead of 36 months;	1.18	1.05 — 1.32	0.0046
Analyzing the change in KCCQ-OSS at 24 months instead of 36 months;	1.17	1.03 — 1.33	0.0149
Using a threshold of $\geq 5$ -point improvement in KCCQ-OSS, rather than any difference, to determine a win	1.23	1.06 — 1.43	0.0079
Performing a similar win ratio analysis in the full TOPCAT trial population	1.12	1.01 — 1.25	0.0261

CI: confidence interval; KCCQ: Kansas City Cardiomyopathy Questionnaire Overall summary score; TOPCAT: Treatment of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist.